

A Universal Scaling Theory for Complexity of Analog Computation

Yaniv S. Avizrats,¹ Shmuel Fishman,¹ and Joshua Feinberg^{2,1}

¹*Department of Physics, Technion, Haifa 32000, Israel.*

²*Department of Physics, University of Haifa at Oranim, Tivon 36006, Israel.*

We discuss the computational complexity of solving linear programming problems by means of an analog computer. The latter is modeled by a dynamical system which converges to the optimal vertex solution. We analyze various probability ensembles of linear programming problems. For each one of these we obtain numerically the probability distribution functions of certain quantities which measure the complexity. Remarkably, in the asymptotic limit of very large problems, each of these probability distribution functions reduces to a universal scaling function, depending on a single scaling variable and independent of the details of its parent probability ensemble. These functions are reminiscent of the scaling functions familiar in the theory of phase transitions. The results reported here extend analytical and numerical results obtained recently for the Gaussian ensemble.

PACS numbers: 5.45-a, 89.79+c, 89.75.D

During the past two decades or so, physicists have been applying methods of statistical physics in studying hard combinatorial optimization problems of computer science [1]. Physical methods, borrowed from statistical physics and the theory of dynamical systems (DS) [2], have been applied recently in studying an easier computer science problem, namely, the computational complexity of an analog computation algorithm which solves linear programming (LP) problems [3, 4]. In this Letter we demonstrate the robustness and universality of the results of [3, 4].

To put things in broader context, we remark that analog computers are ubiquitous computational tools, alongside with their predominating digital counterparts. The most relevant examples of analog computers are VLSI devices implementing neural networks [5], or neuromorphic systems [6], whose structure is directly motivated by the workings of the brain. Various processes taking place in living cells can be considered as analog computation [7] as well.

Linear programming is a P-complete problem [8], i.e. it is representative of all problems that can be solved in polynomial time. The *standard form* of LP is to find

$$\max\{c^T x : x \in \mathbb{R}^n, Ax = b, x \geq 0\} \quad (1)$$

where $c \in \mathbb{R}^n$, $b \in \mathbb{R}^m$, $A \in \mathbb{R}^{m \times n}$ and $m \leq n$. The set generated by the constraints in (1) is a polyhedron. If a bounded optimal solution exists, it is obtained at one of its vertices. The vector defining this optimal vertex can be decomposed (in an appropriate basis) in the form $x = (x_N, x_B)$ where $x_N = 0$ is an $n - m$ component vector, while $x_B = B^{-1}b \geq 0$ is an m component vector, and B is the $m \times m$ matrix whose columns are the columns of A with indices identical to the ones of x_B . Similarly, we decompose $A = (N, B)$, where N is an $(n - m) \times m$ matrix.

A DS whose flow converges to the optimal vertex, introduced by Faybusovich [9], will be studied here. Its

flow $\frac{dx}{dt} = F(x)$ is given in terms of the vector field

$$F(x) = [X - XA^T(AXA^T)^{-1}AX]c, \quad (2)$$

where X is the diagonal matrix $\text{Diag}(x_1 \dots x_n)$. Geometrically, F is the projection of the gradient of the cost function $c^T x$ onto the constraint set, relative to a Riemannian metric, which enforces the positivity constraints $x \geq 0$ [9].

This DS, as it evolves in its continuous phase space in continuous time, models the analog computer in question, which solves the given LP problem. Other dynamical systems are known (also described by ordinary differential equations) that are used to solve various computational problems [9, 10, 11]. Thus, a large set of analytical tools and physical intuition, developed for dynamical systems, turns out to be applicable to the analysis of analog computers.

In contrast, the evolution of a digital computer is described by a dynamical system, discrete both in its phase space and in time. Consequently, the standard theory of computation and computational complexity [8] deals with computation in discrete time and phase space, and is inadequate for the description of analog computers. The analysis of computation by analog devices requires a theory that is valid in continuous time and phase space.

Since the systems in question are physical systems, the computation time is the time required for a system to reach the vicinity of an attractor (a stable fixed point in the present work) combined with the time required to verify that it indeed reached this vicinity. This time is the elapsed time measured by a clock, contrary to standard computation theory, where it is the number of discrete steps.

In our model we assume we have a physical implementation of the flow equation. Thus, the vector field F need not be computed, and the computation time is determined by the convergence time to the attracting fixed point. In other words, the time of flow to the vicinity

of the attractor is a good measure of complexity, namely the computational effort, for the class of continuous dynamical systems introduced above [12].

In this work, following [3, 4] (and in a manner similar to [13, 14]), the complexity will be evaluated probabilistically for an ensemble of LP problems. In this way the worst case scenarios, studied traditionally in computer science, will be ignored, since their probability measure vanishes.

The main result of [3, 4], in which LP problems were drawn from the Gaussian distribution of the parameters of F (namely, the constraints and cost function in (1)), was that the distribution functions of various quantities that characterize the computational complexity, were found to be scaling functions in the limit of LP problems of large size. In particular, it was found that those distribution functions depended on the various parameters only via specific combinations, namely, the scaling variables. Such behavior is analogous to the situation found for the central limit theorem, for critical phenomena [15] and for Anderson localization [16], in spite of the very different nature of these problems. It was demonstrated in [3, 4] how for the implementation of the LP problem on a physical device, methods used in theoretical physics enable to describe the distribution of computation times in a simple and physically transparent form. Based on experience with certain universality properties of rectangular and chiral random matrix models, it was conjectured in [3, 4] that some universality for computational problems should be expected and should be explored. That is, the scaling properties that were found for the Gaussian distributions should hold also for other distributions. In particular, some specific questions were raised in [3, 4]: Is the Gaussian nature of the ensemble unimportant? Are there universality classes [15] of analog computational problems, and if they exist, what are they? In the present work we extend the earlier analysis [3, 4] of the Gaussian distribution to other probability distributions, and demonstrate *numerically* that the distribution functions of various quantities that characterize the computational complexity of the analog computer, which solves LP problems, are indeed *universal* scaling functions. They depend upon the original probability ensemble of inputs only via the scaling variables, that are proportional to the ones found for the Gaussian distribution.

The distribution of constraints and cost function of the LP problems that are used in practice is not known. Therefore, the universality of the distribution functions of the computation time and other quantities related to computational complexity is of great importance. It would imply that distributions found for the idealized systems may be relevant also for the realistic sets of LP problems. In this paper we demonstrate numerically that for several probability distributions of LP problems, the functions of the quantities that characterize the complex-

ity are indeed universal, providing support for the conjecture that universality holds in general.

It was shown in [17] that the flow equations corresponding to (2) are, in fact, part of a system of Hamiltonian equations of motion of a completely integrable system of a Toda type. Therefore, like the Toda system, it is integrable with the formal solution [9]

$$x_i(t) = x_i(0) \exp \left(-\Delta_i t + \sum_{j=1}^m \alpha_{ji} \log \frac{x_{j+n-m}(t)}{x_{j+n-m}(0)} \right) \quad (3)$$

($i = 1, \dots, n-m$), that describes the time evolution of the $n-m$ independent variables $x_{\mathcal{N}}(t)$, in terms of the variables $x_{\mathcal{B}}(t)$. In (3) $x_i(0)$ and $x_{j+n-m}(0)$ are components of the initial condition, $x_{j+n-m}(t)$ are the $x_{\mathcal{B}}$ components of the solution, $\alpha_{ji} = -(B^{-1}N)_{ji}$ is an $m \times (n-m)$ matrix, while

$$\Delta_i = -c_i - \sum_{j=1}^m c_j \alpha_{ji}. \quad (4)$$

For the decomposition $x = (x_{\mathcal{N}}, x_{\mathcal{B}})$ used for the optimal vertex, $\Delta_i \geq 0$ $i = 1, \dots, n-m$, and $x_{\mathcal{N}}(t)$ converges to 0, while $x_{\mathcal{B}}(t)$ converges to $x^* = B^{-1}b$. Note that the analytical solution is only a *formal* one, and does not provide an answer to the LP instance, since the Δ_i depend on the partition of A , and only relative to a partition corresponding to a maximum vertex are all the Δ_i positive.

The second term in (3), when it is positive, is a kind of “barrier”: $\Delta_i t$ must be larger than the barrier before x_i can decrease to zero. In the following we ignore the contribution of the initial condition and denote the value of this term in the infinite time limit by

$$\beta_i = \sum_{j=1}^m \alpha_{ji} \log x_{j+n-m}^*. \quad (5)$$

In order for $x(t)$ to be close to the maximum vertex we must have $x_i(t) < \epsilon$ for $i = 1, \dots, n-m$ for some small positive ϵ , namely $\exp(-\Delta_i t + \beta_i) < \epsilon$, for $i = 1, \dots, n-m$. Therefore we consider

$$T = \max_i \left(\frac{\beta_i}{\Delta_i} + \frac{|\log \epsilon|}{\Delta_i} \right), \quad (6)$$

as the computation time. We denote

$$\Delta_{\min} = \min_i \Delta_i, \quad \beta_{\max} = \max_i \beta_i. \quad (7)$$

The Δ_i can be arbitrarily small when the inputs are real numbers. Such “bad” instances (associated with the possibility of vanishing x_{j+n-m}^* ’s) are rare in the Gaussian probabilistic model of [3, 4], where it was shown that typically $\Delta_{\min} \sim 1/\sqrt{m}$. In this Letter we will show that this scaling behavior of $\Delta_{\min}$ is universal, being valid in a broad class of probability distributions of LP problems.

Consider an ensemble of LP problems in which the components of (A, b, c) are independent identically distributed (i.i.d.) random variables taken from various *even* distributions, with 0 mean and bounded variance. For a probabilistic model of LP instances, $\Delta_{\min}$, $\beta_{\max}$ and T are random variables.

In [3, 4], the components of A , b , and c were taken from the Gaussian distribution (see, e.g., Eqs.(12-18) in [3]) with zero mean and variance σ^2 . It was found analytically, in the large (n, m) limit, that the probability $\mathcal{P}(\Delta_{\min} < \Delta | \Delta_{\min} > 0) \equiv \mathcal{F}^{(n,m)}(\Delta)$ is of the scaling form

$$\mathcal{F}^{(n,m)}(\Delta) = 1 - e^{x_\Delta^2} \operatorname{erfc}(x_\Delta) \equiv \mathcal{F}(x_\Delta) \quad (8)$$

with the scaling variable

$$x_\Delta = a_\Delta (n/m) x'_\Delta, \quad (9)$$

where

$$x'_\Delta = \sqrt{m} \Delta \quad \text{and} \quad a_\Delta (n/m) = \frac{1}{\sqrt{\pi}} \left(\frac{n}{m} - 1 \right) \frac{1}{\sigma}. \quad (10)$$

The scaling function $\mathcal{F}$ contains *all* asymptotic information on Δ . The distribution $\mathcal{F}(x_\Delta)$ is very wide and does not have a variance. Also the average of $1/x_\Delta$ diverges.

The amazing point is that in the limit of large m and n , the probability distributions of $\Delta_{\min}$ depend on the variables m , n and Δ only via the scaling variable x_Δ . If the limit of infinite m and n is taken, so that n/m is fixed, a_Δ is constant. It was verified numerically that for the Gaussian ensemble (8) is a good approximation already for $m = 20$, and $n = 40$.

The existence of scaling functions like (8) for the barrier $\beta_{\max}$ (that is, the maximum of the β_i) defined by (5), and for T defined by (6) (assuming that ϵ is not too small so that the first term in (6) dominates), was verified numerically for the Gaussian distribution [3, 4].

The scaling behavior (8), and similar behavior found for $1/\beta_{\max}$ and $1/T$, rendering their distribution functions $P_\beta(1/\beta)$ and $P_T(1/T)$ scaling functions of the scaling variables x_β and x_T , all associated with the Gaussian ensemble, prompted us, for reasons discussed above, to explore their universality and check their validity for other probability ensembles of LP problems. Thus, we carried numerical calculations of the distribution functions of Δ , β , and $1/T$ for various probability distributions of A , b , and c .

To be specific, we studied: (1) the uniform distribution, in which each entry of (A, b, c) was uniformly distributed between $\pm \frac{1}{2}$; (2) the discrete, bimodal distribution, in which each entry was either $+1$ or -1 with probability $\frac{1}{2}$ each; and finally, (3) the diluted bimodal distribution, in which each entry was either $+1$ or -1 with probability $\frac{p}{2}$ each, or 0 with probability $1 - p$. Here we chose $p = 0.2, 0.5$ in numerical calculations.

For continuous distributions the probability of encountering degenerate solutions (where some of the x_{j+n-m}^* 's in (5) may vanish) is of measure zero, while for the discrete ensembles some regularization was introduced, in order to avoid degenerate situations.

We generated full LP instances (A, b, c) with the probability distribution in question. For each instance the LP problem was solved using the linear programming solver of MatLab. Only instances with a bounded optimal solution were kept, and $\Delta_{\min}$ was computed relative to the optimal partition and optimality was verified by checking that $\Delta_{\min} > 0$. Using the sampled instances we obtain an estimate of

$$\mathcal{F}^{(n,m)}(\Delta) \equiv P_\Delta(\Delta) = \mathcal{P}(\Delta_{\min} < \Delta | \Delta_{\min} > 0) \quad (11)$$

and of the corresponding cumulative distribution functions of the barrier $\beta_{\max}$ and the computation time T .

The solution of the LP problem is used here in order to identify the optimal partition of A into B and N . This enables one to compute $\Delta_{\min}$, $\beta_{\max}$, and T from (6) and (7), and the distribution $P_\Delta(\Delta)$ as well as $P_\beta(1/\beta)$ and $P_T(1/t)$.

For example, the scaling behavior of (11) in the case of the uniform ensemble can be seen in Fig. 1. Similar scal-




FIG. 1: $\mathcal{P}_\Delta$ is plotted as a function of x'_Δ for the uniform distribution for LP problems with $n = 2m$. The number of instances used in the simulation is 121939 for the $m = 20$ case, 91977 for the $m = 30$ case and 112206 for the $m = 40$ case. The number of converging instances for each case is 20000.

ing behavior is found also for P_β and P_T for the uniform distribution. Scaling behavior of this nature was confirmed also for the bimodal distribution, and preliminary results suggest that it holds also for the diluted bimodal distribution.

A natural question which arises is whether the distribution functions P_Δ , P_β and P_T are universal [3, 4]. In other words, do all probability ensembles of LP problems, or at least a large family thereof, yield the same functions P_Δ , P_β and P_T of the scaling variables x_Δ , x_β

and x_T ? We found that the answer to this question is on the affirmative.

Specifically for Δ , plots like Fig. 1 were produced for the variables x'_Δ for the various distributions that were studied. Indeed, these were found to be scaling functions. Then, the scale factors a_Δ , corresponding to (10) were calculated for the various distributions so that P_Δ as a function of x_Δ is the same function for all distributions. (This is done, as usual, by least squares fit.) Indeed, a universal function for P_Δ is found, as is clear from Fig. 2, and it reduces to (8) that was found for the Gaussian ensemble in [3, 4]. It turns out that all the scaling factors that were found in this way for the various distributions are in agreement with (10) (in the way it depends on σ). Similar scaling was found also for the distribution functions P_β and P_T of $1/\beta_{max}$ and $1/T$, respectively.




FIG. 2: P_Δ as a function of x_Δ . Here $n = 2m$ and $m = 40$. The scale factors a_Δ are: $1.044/\sqrt{\pi}$ for the bimodal distribution, $3.603/\sqrt{\pi}$ for the uniform distribution, and $1/\sqrt{\pi}$ for the gaussian distribution. For the Gaussian and bimodal distributions the variance is $\sigma^2 = 1$, while for the uniform distribution $\sigma^2 = 1/12$.

In summary, we find that the asymptotic distribution functions P_Δ , P_β and P_T are the same for all distributions of the parameters studied here, if expressed in terms of the scaling variables x_Δ , x_β and x_T . Furthermore, we find that then the following combinations of scaling factors, σa_Δ , a_β and σa_T are independent of the probability distribution. Therefore, in particular, a_Δ should satisfy (10) that was found for the Gaussian distribution, while a_β and a_T are yet to be found analytically.

Based the results presented in this paper, as well as the results of [3, 4], we conjecture that the scaling behavior of the various distribution functions (and the corresponding scaling factor) is *universal*, i.e., that it is robust and

should be valid in a large class of ensembles of LP problems, in which the (A, b, c) data are taken from even distributions, with zero mean and finite variance.

It is a pleasure to thank our colleagues Asa Ben-Hur and Hava Siegelmann for advice and discussions. This research was supported in part by the Shlomo Kaplan-sky Academic Chair, by the Technion-Haifa University Collaboration Fund, by the US-Israel Binational Science Foundation (BSF), by the Israeli Science Foundation, and by the Minerva Center of Nonlinear Physics of Complex Systems.

-
- [1] M. Mézard, G. Parisi, and M.-A. Virasoro. *Spin Glass Theory and Beyond*. World Scientific, Singapore, 1993.
 - [2] E. Ott, *Chaos in Dynamical Systems*. Cambridge University Press, Cambridge, England, 1993.
 - [3] A. Ben-Hur, J. Feinberg, S. Fishman, and H.T. Siegelmann, *J. Complexity* **19**, 474 (2003) (cs.CC/0110056).
 - [4] A. Ben-Hur, J. Feinberg, S. Fishman, and H.T. Siegelmann, *Phys. Lett. A* **323**, 204 (2004) (cond-mat/0110655).
 - [5] J. Hertz, A. Krogh, and R. Palmer. *Introduction to the Theory of Neural Computation*. Addison-Wesley, Redwood City, 1991.
 - [6] C. Mead. *Analog VLSI and Neural Systems*. Addison-Wesley, 1989.
 - [7] D. Bray. *Nature* **376**, 307 (1995); A. Ben-Hur and H.T. Siegelmann. *Proceedings of MCU 2001, Lecture Notes in Computer Science* **2055**, pages 11-24, M. Margenstern and Y. Rogozhin (Editors), Springer Verlag, Berlin 2001 (and references therein).
 - [8] C. Papadimitriou. *Computational Complexity*. Addison-Wesley, Reading, Mass., 1995.
 - [9] L. Faybusovich. *IMA Journal of Mathematical Control and Information* **8**, 135 (1991).
 - [10] R. W. Brockett. *Linear Algebra and Its Applications* **146**, 79 (1991); M.S. Branicky. Analog computation with continuous ODEs. In *Proceedings of the IEEE Workshop on Physics and Computation*, pages 265–274, Dallas, TX, 1994.
 - [11] U. Helmke and J.B. Moore. *Optimization and Dynamical Systems*. Springer Verlag, London, 1994.
 - [12] H.T. Siegelmann, A. Ben-Hur and S. Fishman, *Phys. Rev. Lett.* **83**, 1463 (1999); A. Ben-Hur, H.T. Siegelmann, and S. Fishman. *J. Complexity* **18**, 51 (2002).
 - [13] S. Smale. *Math. Programming* **27**, 241 (1983). M.J. Todd. *Mathematics of Operations Research* **16**, 671 (1991).
 - [14] R. Shamir. *Management Science* **33**(3), 301 (1987).
 - [15] K.G. Wilson and J. Kogut, *Phys. Rep.* **12**, (1974) 75.
 - [16] E. Abrahams, P. W. Anderson, D. C. Licciardello and T. V. Ramakrishnan, *Phys. Rev. Lett.* **42** (1979) 673.
 - [17] L. Faybusovich. *Physica* **D53**, 217 (1991).